

44. Method of Descent

In this exercise we will apply the method of descent to solve the wave equation on \mathbb{R}^2 for a particular set of initial conditions. The idea is to help you understand the key ideas and notation of the method. It is a combination of results from Sections 5.1–4.

Consider the wave equation on \mathbb{R}^2 with initial conditions

$$\begin{aligned} \partial_t^2 u - \Delta u &= 0 \text{ on } (x, t) \in \mathbb{R}^2 \times (0, \infty), \\ u(x, 0) &= g(x) = \chi_{[0, \infty)}(x_1), \quad \partial_t u(x, 0) = h(x) = 0. \end{aligned}$$

- (a) In Section 5.4 we define $\bar{u} : \mathbb{R}^3 \times [0, \infty) \rightarrow \mathbb{R}$ associated to the function u . Explain why $\bar{g}(x_1, x_2, x_3) = \chi_{[0, \infty)}(x_1)$ and $\bar{h} = 0$ (or give the definition of bar). Why does \bar{u} solve the wave equation on \mathbb{R}^3 ? (note, the Laplacians are different in different dimensions). (2 Points)
- (b) Conversely, prove that a solution \bar{u} to the 3-dimensional wave equation that does not depend on x_3 gives a solution to the 2-dimensional wave equation. (2 Points)
- (c) By (a) and (b), we now must solve a wave equation on \mathbb{R}^3 . The key to solving the 3-dimensional wave equation is to consider the (spatial-)spherical means

$$U(x, t, r) = \frac{1}{4\pi r^2} \int_{\partial B(x, r)} \bar{u}(z, t) \, d\sigma(z),$$

and likewise let G and H be the spherical means of \bar{g} and \bar{h} respectively. Show that

$$G(x, r) = \begin{cases} 0 & \text{for } x_1 \leq -r \\ \frac{1}{2} \frac{x_1 + r}{r} & \text{for } |x_1| \leq r \\ 1 & \text{for } r \leq x_1 \end{cases} \quad \text{and } H(x, r) = 0.$$

You may use the following geometric fact: for $-R < a < b < R$, the surface area of the part of the sphere $\partial B(0, R)$ with $a < x_1 < b$ is $2\pi R(b - a)$. (4 Points)

- (d) We know by Lemma 5.2 that U obeys the Euler-Poisson-Darboux equation. Let $\tilde{U}(x, t, r) := rU(x, t, r)$. Show that \tilde{U} obeys the following PDE

$$\begin{aligned} \partial_t^2 \tilde{U} - \partial_r^2 \tilde{U} &= 0 \text{ on } (t, r) \in [0, \infty) \times [0, \infty), \\ \tilde{U}(x, 0, r) &= rG(x, r), \quad \partial_t \tilde{U}(x, 0, r) = rH(x, r). \end{aligned}$$

Note that x plays no role in this PDE, so we can think of it as a family of PDEs parametrised by x . (2 Points)

- (e) Thus we see that \tilde{U} obeys the 1-dimensional wave equation on the *half-line* $r \in [0, \infty)$. This is solved by a trick using reflection, and the formula is at the end of Section 5.1. We only

need the solution for small r , so it is enough to consider the case $0 \leq r \leq t$. In this case, show

$$\tilde{U}(x, t, r) = \begin{cases} 0 & \text{for } x_1 \leq -(t+r) \\ \frac{1}{4}(x_1 + t + r) & \text{for } -(t+r) \leq x_1 \leq -(t-r) \\ \frac{1}{2}r & \text{for } |x_1| \leq t-r \\ \frac{1}{4}(x_1 - t + 3r) & \text{for } t-r \leq x_1 \leq t+r \\ r & \text{for } x_1 \geq t+r. \end{cases}$$

(4 Points)

- (f) Recover \bar{u} from \tilde{U} using a certain property of spherical means. (1 Point + 2 Bonus Points)

Observe that \bar{u} does not depend on x_3 . So by part (b) we have a solution to the 2-dimensional wave equation:

$$u(x_1, x_2, t) = \begin{cases} 0 & \text{for } x_1 < -t \\ 0.25 & \text{for } x_1 = -t \\ 0.5 & \text{for } -t < x_1 < t \\ 0.75 & \text{for } x_1 = t \\ 1 & \text{for } x_1 > t. \end{cases}$$

This solution has jump discontinuities, but this is unsurprising since the initial conditions also had them.

Solution.

- (a) Recall the definition $\bar{u}(x_1, x_2, x_3, t) = u(x_1, x_2, t)$. In words, the bar function is the same formula but considered in a higher dimensional space. This explains \bar{g} and \bar{h} . So it is constant in the x_3 dimension and $\partial_3 \bar{u} = 0$.

If we write out the wave equation on \mathbb{R}^3 fully

$$(\partial_t^2 - \partial_1^2 - \partial_2^2 - \partial_3^2)\bar{u} = (\partial_t^2 - \partial_1^2 - \partial_2^2)u - \partial_3^2 \bar{u} = 0.$$

- (b) By the same reasoning

$$(\partial_t^2 - \partial_1^2 - \partial_2^2)\bar{u} = \partial_3^2 \bar{u} = 0.$$

Hence we get the solution $u(x_1, x_2, t) = \bar{u}(x_1, x_2, 0, t)$. The choice $x_3 = 0$ is not significant, because \bar{u} is constant in x_3 . Any other choice gives the same thing.

- (c) By definition

$$G(x, r) = \frac{1}{4\pi r^2} \int_{\partial B_3(x, r)} \chi_{[0, \infty)}(z_1) d\sigma(z).$$

We use B_3 here to make clear this is a ball in 3-dimensional space. If this ball lies entirely in the space with $z_1 \geq 0$ then the integrand is always 1 and the integral is just the surface area of the sphere. This occurs if x_1 (the first coordinate of the centre of the ball) is greater than the radius r . Likewise, if $x_1 < -r$ then the integrand is always zero.

So it remains to handle the case $-r \leq x \leq r$. The integral is

$$G(x, r) = \frac{1}{4\pi r^2} \int_{\partial B_3(x, r) \cap \{0 \leq x_1 \leq r\}} 1 \, d\sigma(z) = \frac{1}{4\pi r^2} \times 2\pi r(r + x_1) = \frac{1}{2} \frac{x_1 + r}{r}.$$

(d) We just differentiate

$$\partial_t^2(rU) - \partial_r^2(rU) = r\partial_t^2 U - \partial_r(U + r\partial_r U) = r\partial_t^2 U - 2\partial_r U - r\partial_r^2 U.$$

This is r multiplied by the Euler-Poisson-Darboux equation for $n = 3$. Since U solves this equation, we get 0 on the right hand side. For the initial conditions $\tilde{U}(x, 0, r) = rU(x, 0, r) = rG(x, r)$ and likewise for H .

(e) The solution of the wave equation on the half line, for $0 \leq r \leq t$ is

$$\begin{aligned} \tilde{U}(x, t, r) &= \frac{1}{2} [\tilde{G}(x, t+r) - \tilde{G}(x, t-r)] + \frac{1}{2} \int_{t-r}^{t+r} \tilde{H}(x, y) \, dy \\ &= \frac{1}{2} [(t+r)G(x, t+r) - (t-r)G(x, t-r)] + \frac{1}{2} \int_{t-r}^{t+r} 0 \, dy, \end{aligned}$$

since $\tilde{H} = rH = 0$. Under the assumption $0 \leq r \leq t$, you can see that $-(t+r) \leq -(t-r) \leq 0 \leq (t-r) \leq (t+r)$. Thus there are five cases intervals to consider

$$\tilde{U}(x, t, r) = \begin{cases} \text{for } x_1 \leq -(t+r) : & 0 - 0 \\ \text{for } -(t+r) \leq x_1 \leq -(t-r) : & \frac{1}{4}(x_1 + t+r) - 0 \\ \text{for } |x_1| \leq t-r : & \frac{1}{4}(x_1 + t+r) - \frac{1}{4}(x_1 + t-r) \\ \text{for } t-r \leq x_1 \leq t+r : & \frac{1}{2}(t+r) - \frac{1}{4}(x_1 + t-r) \\ \text{for } x_1 \geq t+r : & \frac{1}{2}(t+r) - \frac{1}{2}(t-r) \end{cases}$$

(f) We know that a function is equal to the limit of its spherical mean as the radius goes to zero, $\bar{u}(x, t) = \lim_{r \rightarrow 0} U(x, t, r) = \lim_{r \rightarrow 0} r^{-1} \tilde{U}(x, t, r)$. The first, third, and fifth cases of \tilde{U} give

$$\bar{u}(x_1, x_2, x_3, t) = \begin{cases} 0 & \text{for } x_1 < -t \\ \frac{1}{2} & \text{for } -t < x_1 < t \\ 1 & \text{for } x_1 > t. \end{cases}$$

To find the value of $\bar{u}(x, t)$ for $x_1 = -t$, we see that this sits in the second case of \tilde{U} . We get

$$\lim_{r \rightarrow 0} \frac{1}{4} \frac{x_1 + t + r}{r} = \lim_{r \rightarrow 0} \frac{1}{4} \frac{0 + r}{r} = \frac{1}{4}.$$

Similarly $\bar{u}(t, x_2, x_3, t) = \frac{3}{4}$.

This behaviour at the jump discontinuities is typical for these averaging methods, it gives the value at the jump as the average of the two sides of the discontinuity. Properly, when we have functions which are not twice continuously differentiable we should use distributions and weak solutions. In that context, the value of the function at the jump is not significant, it is just an artifact of using spherical means.

45. Plane Waves.

Suppose that $u : \mathbb{R}^n \times \mathbb{R} \rightarrow \mathbb{R}$ is a solution to the following modified wave equation:

$$\frac{\partial^2 u}{\partial t^2} - \sum_{j=1}^n c_j^2 \frac{\partial^2 u}{\partial x_j^2} = 0, \quad (*)$$

where $c_1, \dots, c_n > 0$ are constants.

- (a) Let $\alpha \in \mathbb{R}^n$ be a unit vector $\|\alpha\| = 1$, $\mu \in \mathbb{R}$ and $F : \mathbb{R} \rightarrow \mathbb{R}$ a twice continuously differentiable function. Show that

$$u(x, t) := F(\alpha \cdot x - \mu t)$$

is a solution of (*) exactly when

$$\mu^2 = \sum_{j=1}^n \alpha_j^2 c_j^2$$

or F is linear. Solutions of (*) with this form are called *plane waves*. (2 Points)

- (b) For the solutions in (a), examine whether the following property holds for all $x \in \mathbb{R}^n$ and $t \in \mathbb{R}$:

$$u(x, t) = u(x - \mu t \alpha, 0).$$

Interpret this equation in terms of direction and speed. (3 Points)

Solution.

- (a) We apply the chain rule to differentiate F :

$$\frac{\partial^2 u}{\partial t^2} - \sum_{j=1}^n c_j^2 \frac{\partial^2 u}{\partial x_j^2} = (-\mu)^2 F'' - \sum_{j=1}^n c_j^2 (\alpha_j)^2 F'' = \left(\mu^2 - \sum_{j=1}^n c_j^2 \alpha_j^2 \right) F''.$$

Clearly this is zero only if the relation between μ and α holds or if $F'' = 0$.

- (b) This property does hold, because of the normalisation condition $|\alpha| = \alpha \cdot \alpha = 1$:

$$u(x, t) = F(\alpha \cdot x - \mu t) = F(\alpha \cdot (x - \mu t \alpha)) = F(\alpha \cdot (x - \mu t \alpha) - \mu t) = u(x - \mu t \alpha, 0).$$

This shows that plane waves are constant along the planes $x \cdot \alpha = \text{const.}$. If we consider a line parallel, then the problem is reduced to the one dimensional wave equation with speed μ . Hence we say the wave is moving in the direction α .

There are other sorts basic waves; spherical waves and standing waves are two important examples. In three dimensions, if a solution only depends on $r = |x|$ then the wave equation becomes

$$0 = \partial_t^2 u - \partial_r^2 u - \frac{2}{r} \partial_r u = \frac{1}{r} (\partial_t^2 - \partial_r^2)(ru).$$

This is again a one dimensional wave equation, solved by $u(r, t) = r^{-1}F(r - t) + r^{-1}G(r + t)$. The interpretation here is that there are inward and outward moving spheres, but the amplitude is diminished/concentrated as the radius is changed.

A standing wave is one whose peaks do not move in space, it only oscillates in time. Simple standing waves separate into the form $u(x, t) = \tilde{u}(x) \sin(\omega t)$. The profile of the wave (the \tilde{u} part) is governed by the equation

$$0 = (\partial_{tt} - \Delta)u = (-\omega^2\tilde{u} - \Delta\tilde{u}) \sin(\omega t).$$

Alternatively, this arises from taking the Fourier transformation in t , namely $\hat{u}(x, \omega) = \int u(x, t)e^{-i\omega t} dt$, and considering solutions with a constant frequency ω .

46. Electromagnetic Waves.

In physics, electrical and magnetic fields are modelled as time-dependent vector fields, which mathematically are smooth functions $E, B : \mathbb{R}^3 \times \mathbb{R} \rightarrow \mathbb{R}^3$. Through a series of experiments in the 18th and 19th centuries, the existence and properties of these fields were discovered. Importantly, it was discovered that the two phenomena were connected (both magnets and static electricity had been known since antiquity). In 1861 James Clerk Maxwell published a series of papers summarising electromagnetic theory, including a collection of 20 differential equations. Over time these were further reduced to the following four (by Heaviside 1884 using vector notation), called *Maxwell's Equations*:

$$\begin{aligned} \nabla \cdot E &= \frac{1}{\varepsilon_0} \rho & \nabla \times E &= -\frac{\partial B}{\partial t} \\ \nabla \cdot B &= 0 & \nabla \times B &= \mu_0 J + \varepsilon_0 \mu_0 \frac{\partial E}{\partial t}. \end{aligned}$$

As is usual, the ∇ operator acts on the spatial coordinates x , and the \times denotes the cross product of \mathbb{R}^3 . The constants ε_0 , the electrical permittivity, and μ_0 , the magnetic permeability, are approximately $\varepsilon_0 \approx 8,854 \cdot 10^{-12} \frac{\text{A}\cdot\text{s}}{\text{V}\cdot\text{m}}$ and $\mu_0 \approx 1,257 \cdot 10^{-6} \frac{\text{V}\cdot\text{s}}{\text{A}\cdot\text{m}}$ (V=Volt, s=Seconds, A=Ampere and m=Metre) in a vacuum. Electrical charges are included via the charge density ρ and electric currents are the movements of charges, $J := v\rho$ for a velocity field v .

The two equations with divergence were formulated by Gauss, the curl of the electric field is due to Faraday, and the curl of the magnetic field is due to Ampère. The last term in Ampère's law that has the time-derivative of the electrical field was an addition of Maxwell. With this correction, he was able to derive the equations for electromagnetic waves, as you will now do.

(a) Let E and B be solutions to Maxwell's equations in the absence of electric charges, $\rho = 0, J = 0$. Show that they each satisfy a modified wave equation (Question 41). You may use without proof the identity $\nabla \times (\nabla \times f) = \nabla(\nabla \cdot f) - \Delta f$ for smooth functions $f : \mathbb{R}^3 \rightarrow \mathbb{R}^3$.

(3 Bonus Points)

(b) Predict the speed of these waves.

(2 Bonus Points)

- (c) Argue that Ampère's law in its original form $\nabla \times B = \mu_0 J$ violates the conservation of charge ρ under some conditions. Refer to Exercise Sheet 5 Question 16 for the definition of a conservation law. Thereby derive Maxwell's additional term. *(3 Bonus Points)*

Solution.

- (a) Suppose we have solutions E, B . As suggested by the hint, we take the curl of the curl equations. Because curl is a linear operator (and derivatives commute) we may write

$$\nabla \times \nabla \times E = -\frac{\partial}{\partial t} \nabla \times B = -\frac{\partial}{\partial t} \varepsilon_0 \mu_0 \frac{\partial E}{\partial t} = -\varepsilon_0 \mu_0 \frac{\partial^2 E}{\partial t^2}.$$

On the other hand, we know the twice curl of E is $\nabla(\nabla \cdot E) - \Delta E = \nabla(0) - \Delta E$, using Gauss' law of electric fields. Rearranging we get a modified wave equation:

$$\frac{\partial^2 E}{\partial t^2} = \frac{1}{\varepsilon_0 \mu_0}$$

and likewise for B .

- (b) We expect that the speed is given by μ as in Question 41(b), and this can be calculated from the coefficients c_j and the direction α . In this case, the coefficients are the same in each coordinate direction, so factor out:

$$\mu = \sqrt{\sum a_j^2 c^2} = c|\alpha| = c = \frac{1}{\sqrt{\varepsilon_0 \mu_0}} \approx 299\,800\,000 \text{ ms}^{-1}.$$

This is the speed of light. The speed of light had first been calculated nearly 200 years earlier by Romer using astronomical observations of Jupiter and its moons, and would in 1862 measured with less than 1% error. The electrical constant had been determined only 5 years earlier with experiments with capacitors by Weber and Kohlrausch. The magnetic constant is fixed by the definition to be $4\pi \cdot 10^{-7}$. The measurements were good enough in Maxwell's day to see that these were close, and on this basis Maxwell hypothesised light was an electromagnetic wave.

- (c) We saw in Question 12 that a quantity, be it mass or in this case electrical charge, is conserved when the change of density is equal to the negative of the divergence of the flow (using the divergence theorem, the divergence of the flow is the amount of substance leaving a small ball around that point). Symbolically, $\partial_t \rho = -\nabla \cdot (v\rho) = \nabla \cdot J$. If we take the divergence of Ampère's version we have

$$0 = \nabla \cdot (\nabla \times B) = \mu_0 \nabla \cdot J.$$

This is only true when the charge density ρ is constant. As Ampère's experiment used two wires with constant currents, this was true in his experiment.

But in general we should add another term $\nabla \times B = \mu_0 J + G$. Applying the divergence now, we see that

$$\nabla \cdot G = -\mu_0 \nabla \cdot J = \mu_0 \frac{\partial}{\partial t} \rho = \mu_0 \varepsilon_0 \frac{\partial}{\partial t} \nabla \cdot E.$$

Hence we conclude that $G = \mu_0 \varepsilon_0 \partial_t E + \nabla \times g$. Taking the simplest possibility, $g = 0$, gives Maxwell's correction.

Note that we shows that each component of the electric and magnetic fields solve the wave equation, but this is a necessary condition. Faraday's law show that there is a dependence between the two fields. And both fields must have zero divergence, which creates a dependence directly between the components. For example, consider if all of E_i are plane waves travelling in the x_3 direction, so E depends only on x_3 . Then $\nabla \cdot E = 0$ implies $E_3 = 0$. The relations between the components is polarization. For example, a solution such as $E_1 = E_1(x_3 - ct)$, $E_2 = E_3 = 0$ is a wave travelling in the x_3 direction, but polarized in the x_1 direction.

Solutions are due on Tuesday 12 noon, the day before the tutorial. Please email to r.ogilvie@math.uni-mannheim.de. One possibility is to write your solutions neatly by hand and then scan them with your phone to make a pdf. There are many apps that do this; two examples on Android are 'Tiny Scanner' and 'Simple Scanner'.