

77. Orientable hypersurfaces defined by an equation.

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ a smooth map, $q \in \mathbb{R}$ a point in its range so $X := f^{-1}(\{q\}) \neq \emptyset$, and f is submersive at all points $x \in X$. Show that X is an $(n - 1)$ -dimensional orientable submanifold.

Hint. Let ω be the volume form on \mathbb{R}^n and F the gradient field of f (i.e $T_x(f)(v) = F(x) \cdot v$). Investigate $i_F \omega|_X$, defined in Definition 3.11.

In Class Exercises

78. Orientable manifolds.

- (a) Show that the n -dimensional sphere \mathbb{S}^n is orientable by finding an oriented atlas.
Hint. For the sphere, consider the atlas that uses stereographic projection. An extra trick is also needed.
- (b) Show that the Möbius band is not orientable.
Hint. This is difficult. Good luck.
- (c) Let X and Y be orientable manifolds. Show that the Cartesian product $X \times Y$ is also orientable.
- (d) Let X be a manifold. Show that every coordinate neighbourhood of X is orientable. More precisely, let (U, ϕ) be a chart of X with $\phi = (\phi_1, \dots, \phi_n) : U \rightarrow \mathbb{R}^n$, and show that $d\phi_1 \wedge \dots \wedge d\phi_n$ is a non-vanishing n -form on U .
- (e) Prove that the tangent bundle of any manifold is orientable.

79. Integration on \mathbb{R} .

Consider the following integration by substitution with $f(x) = x^2$:

$$\int_{-1}^2 2xe^{-x^2} dx = \int_{-1}^2 e^{f(x)} f'(x) dx = \int_{f(-1)}^{f(2)} e^u du = \int_1^4 e^u du.$$

This seems very similar to Corollary 3.22, except that f is not a diffeomorphism! Also consider the substitution with $v = -x$

$$\int_{-1}^2 e^{-x} dx = \int_1^{-2} e^u (-du) = \int_{-2}^1 e^u du.$$

The aim of this exercise is to understand why this all works for \mathbb{R} .

First, extend Corollary 3.22 for orientation-reversing diffeomorphisms.

Second, define

$$\int_a^b g(x) dx = \text{sign}(b - a) \int_{\mathbb{R}} \chi_{[a,b]}(x)g(x) dx.$$

Show that it obeys the rule $\int_a^b + \int_b^c = \int_a^c$.

Third, let $f : [a, b] \rightarrow \mathbb{R}$ be a continuously differentiable function. The set $(f')^{-1}[\{0\}]$ is closed, and therefore the countable union of closed intervals J_i (single points are considered as the interval $[x, x]$). Taking the endpoints of these intervals gives a partition $\{t_i\}$ of $[a, b]$ such that on each interval (t_i, t_{i+1}) the function f' is either non-zero or identically zero. Prove

$$\int_a^b g(f(x))f'(x) dx = \int_{f(a)}^{f(b)} g(u) du.$$