

**7. Royale with Cheese**

Recall Burgers' equation from Example 1.5 of the lecture script:

$$\dot{u} + u\partial_x u = 0,$$

for  $u : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ . In this question we will apply the method of characteristics to solve this equation for the initial condition  $g(x) = 2x$ .

- (a) According to Theorem 1.4, there is a unique  $C^1$  solution to this initial value problem, at least when  $t$  is small. For how long does the theorem guarantee that the solution exists uniquely? *(1 point)*
- (b) Suppose that  $u$  is a solution to this equation and suppose that  $(x(s), t(s))$  is a path in the domain of  $u$ . What is the  $s$  derivative of  $u$  along this path? What constraints should we place on the derivatives of  $x$  and  $t$ ? *(2 points)*
- (c) On an  $(x, t)$ -plane draw the characteristics. *(1 point)*
- (d) Finally, derive the following solution to the initial value problem: *(2 points)*

$$u(x, t) = \frac{2x}{1 + 2t}.$$

- 8. Linear Partial Differential Equations** Consider a PDE of the form  $F(\nabla u(x), u(x), x) = 0$ . Suppose that  $F$  is linear in the derivatives and has continuously differentiable coefficients. That is, it can be written in the form

$$F(p, z, x) = b(z, x) \cdot p + c(z, x)$$

with  $b$  and  $c$  continuously differentiable. Show that the characteristic curves  $(x(s), z(s))$  for  $z(s) := u(x(s))$  can be described by ODEs that are independent of  $p(s) := \nabla u(x(s))$ . *(4 points)*

- 9. Solving PDEs** Solve the initial value problems of the following PDEs using the method of characteristics. You may assume that  $g$  is continuously differentiable on the corresponding domain.

- (a)  $x_2\partial_1 u - x_1\partial_2 u = u$  on the domain  $x_1, x_2 > 0$ , with initial condition  $u(x_1, 0) = g(x_1)$ . *(3 points)*
- (b)  $x_1\partial_1 u + 3x_2\partial_2 u + \partial_3 u = 2u$  on  $x_1, x_2 \in \mathbb{R}, x_3 > 0$ , with initial condition  $u(x_1, x_2, 0) = g(x_1, x_2)$ . *(3 points)*
- (c)  $u\partial_1 u + \partial_2 u = 1$  on the domain  $x_1, x_2 > 0$ , with initial condition  $u(x_1, x_1) = \frac{1}{2}x_1$ . *(4 points)*