

17. Convolved

The convolution of two functions $f, g : \mathbb{R}^n \rightarrow \mathbb{R}$ is defined by

$$(f * g)(x) := \int_{\mathbb{R}^n} f(y)g(x-y) dy.$$

- (a) Let $f_n(x) = 0.5n$ for $x \in [-n^{-1}, n^{-1}]$ and 0 otherwise. Show that the following bounds hold

$$\inf_{|y| \leq n^{-1}} g(y) \leq (g * f_n)(0) \leq \sup_{|y| \leq n^{-1}} g(y).$$

(2 points)

- (b) Suppose now that g is continuous. Show that $(g * f_n)(0) \rightarrow g(0)$ as $n \rightarrow \infty$. (2 points)

- (c) Show that the convolution of C_0^∞ -functions on $\mathbb{R}^n$ is a bilinear, commutative, and associative operation. (1+2+2 Points)

18. Distributions

- (a) Choose any compact set $K \subset \mathbb{R}$. Since it is bounded, there exists $R > 0$ with $K \subseteq [-R, R]$. Now choose any test function $\phi \in C_0^\infty(\mathbb{R})$ with compact support in K . Since it is continuous, $\sup_{x \in K} |\phi(x)|$ is finite. Prove the following inequality (1 point)

$$\left| \int_0^\infty \phi(x) dx \right| \leq 2R \sup_{x \in K} |\phi(x)|.$$

- (b) Define the Heaviside function $H : \mathbb{R} \rightarrow \mathbb{R}$ by $H(x) := 1$ for $x \geq 0$ and $H(x) := 0$ for $x < 0$. Show that the distribution associated to the Heaviside function

$$F_H : C_0^\infty(\mathbb{R}) \rightarrow \mathbb{R}, \phi \mapsto \int_0^\infty \phi(x) dx$$

is in fact a distribution on $\mathbb{R}$ using part (a) and Definition 2.14 directly. (1 point)

- (c) Calculate the first and second derivatives of H as a distribution. If they are regular distributions, describe the corresponding function. (2 points)

- (d) Consider the circle $C = \{x^2 + y^2 = 1\} \subset \mathbb{R}^2$. Show that

$$G(\varphi) := \int_C \varphi d\sigma$$

defines a distribution in $\mathcal{D}'(\mathbb{R}^2)$. Note that the $d\sigma$ indicates this is an integration over the submanifold C . Does there exist a locally integrable function $g : \mathbb{R}^2 \rightarrow \mathbb{R}$ with

$$G(\varphi) = \int_{\mathbb{R}^2} g \varphi dx$$

for all $\varphi \in C_0^\infty(\mathbb{R})$? (Hint. Use Lemma 2.15) (2 Points + 2 Bonus Points)

19. Delta Quadrant

- (a) Prove that the support of the delta distribution δ is $\{0\}$. (2 points)

- (b) Argue from Lemma 2.12 that δ is the limit of the standard mollifier $(\lambda_\epsilon)_{\epsilon>0}$ as $\epsilon \downarrow 0$ as a sequence of distributions. (1 point)

- (c) Prove for any distribution F that the convolution with δ is again F . (2 points)